Инструкция по применению для одноразовых игл гиподермальных TRO-CAN

1. Проверьте целостность индивидуальной упаковки.
2. Перед открытием блистера убедитесь, что защитный колпачок иглы на месте, затем снимите его.
3. Иглу следует использовать сразу после открытия упаковки.
4. Утилизируйте изделие после однократного использования в специальный контейнер для острых предметов.
5. Не используйте изделие, если упаковка повреждена или вскрыта.
6. Не используйте изделие с истекшим сроком годности.

/Круглая печать: Отдел контроля и обеспечения качества, ТРОГЕ МЕДИКАЛ ГМБХ (TROGE MEDICAL GMBH), ТРОГЕ Гамбург/

Я, переводчик Бурак Анастасия Михайловна, подтверждаю достоверность совершенного мною перевода текста документа с английского языка на русский язык.

Регистрация МИ в Росздравнадзоре
www.nevacert.ru | info@nevacert.ru

TROGE MEDICAL GMBH

TROGE
HAMBURG

Instruction for Use
for
TRO-CAN Hypodermic Needles for single use

1. Verify individual package integrity.
2. Make sure, that the protector is in place before opening the blister.
3. The needle should be used immediately after opening.
4. Discard in approved sharps collector after single use.
5. Do not use if package is damaged or opened.
6. Do not use expired goods.

Department for Quality Control and Quality Assurance
TROGE
HAMBURG
TROGE MEDICAL GMBH

Регистрация МИ в Росздравнадзоре
www.nevacert.ru | info@nevacert.ru